

Consanguinity and its effect on fertility and infant and child mortality in Egypt

By

Rita G. Khayat and Prem C. Saxena

Abstract

This paper examines the effect of consanguineous marriages on fertility and infant and child mortality in Egypt using country's Demographic Health Survey 2000 data - a nationally representative sample of 16957 households from six governorates of Egypt that includes 15573 ever-married women aged 15-49. These women have been grouped into three separate categories of marriages, namely, 'close consanguineous', 'remote consanguineous' and 'non-consanguineous'. GLIM and logistic regression models have been used to see the impact of consanguinity on fertility and offspring's mortality, respectively after exercising statistical controls on selected socio-economic variables. The results show higher fertility among close consanguineous and remote consanguineous couples. The risk of infant mortality was higher by 30% and 19% in these two groups of women, respectively. Similarly, the risk of child mortality is found elevated among the close consanguineous couples by more than 50% and among remote consanguineous couples by 27% as compared to non-consanguineous unions.

Introduction

Consanguinity, defined as the marriage or unions between individuals of the same blood, is widely practiced in countries of Asia and Africa especially in societies where Islam prevails but its prevalence is low in Western countries. High rates of these marriages are reported in most of the Arab countries. For example, in Sudan 65% of women are married to a relative (Federal Ministry of Health, 1995), Saudi Arabia 57.7% (El Hamzi et al., 1995), Jordan 51.3% (Khoury and Massad, 1992), United Arab Emirates 50.5% (Al-Gazali et al., 1997), Tunisia 49% (Ministry of Public Health, 1996), Egypt 40% (National Population Council, 1996), Yemen 40% (Jurdi and Saxena, 2003) and in Kuwait 36% (Ministry of Health, 1996).

The practice of consanguinity is very old; it was common between the ancient Egyptians of the reigning dynasty to keep the "royal blood" (Stern, 1949). It is known that their offspring have an increased risk of morbidity and mortality. This is because, genetically, they might inherit autosomal recessive conditions or congenital malformations that appear at time of birth or later in childhood. In fact, single gene disorders is common in Eastern Mediterranean families due to the

practice of consanguinity that tends to retain rare mutations within affected families, who may contain a high frequency of mutation carriers. Genetic disorders and congenital abnormalities occur in about 2%-5% of all live births, account for up to 30% of pediatric hospital admissions and cause about 50% of childhood deaths in industrialized countries (Emery and Rimoin, 1990). They are the leading cause of infant mortality in United Arab Emirates (Ministry of health, 1992) and the second leading cause in Bahrain, Kuwait, Oman and Qatar (Ministry of Health of Bahrain, 1991; WHO, 1993; Ministry of Public Health of Qatar, 1993). In Saudi Arabia, 25%-35% of perinatal deaths in two hospitals were attributed to congenital malformations (Agwiser, 1990; Haque and Bashi, 1988). Thus, congenital and genetically determined disorders are very common in the Eastern Mediterranean region. The prevailing high rate of consanguineous marriages is considered to be the root cause of these problems. In a study of 3000 consecutive neonates delivered in a maternity hospital in Giza, Egypt, who were subjected to full clinical and genetic evaluation, parental consanguinity was found in 31.8% of all cases and in 55% of malformed cases, thus illustrating the effects of consanguinity (Temtamy et al., 1998). Recently, the *Jordan Times* dated Monday, March 13, 2000, reported a newly discovered hereditary neurological disorder identified in four Jordanian villages, that causes partial paralysis, has been attributed to the practice of consanguinity. It is believed that off-springs of consanguineous couples are born with biological weaknesses that render them less resistant to infant and childhood diseases resulting in their high mortality.

Consanguinity has been reported as the most important cause of genetically associated mortality in developing countries (Guo, 1993). Couples who are related by blood are more likely to experience the death of an offspring than those not related by blood (Bittles, 1994). A study in Pakistan, found that first cousin marriages are 1.18 times more likely to experience a child's death by its fifth birthday than couples not related by blood (Shah et al., 1998). Another study demonstrated a 3-fold increase of postneonatal mortality and childhood morbidity in the off-springs of couples having consanguineous marriages. It was estimated that 60% of the mortality and morbidity of children, in Pakistan, could be eliminated if these marriages are stopped (Bunday and Alam, 1993). Of late, Banerjee and Roy (2002) found a linkage between the genetic effect of consanguinity and offspring mortality among close consanguineous marriages in India after controlling for the non-genetic determinants. However, the relationship between consanguinity and offspring mortality is still complex and unclear due to the difficulty in classifying death as genetic or non-genetic. In fact, the role of the latter factors has always been an impediment to the understanding of the possible linkage between parental consanguinity and offspring mortality. Among non-genetic factors, socioeconomic status, housing conditions and quality of sanitation are closely associated with infant and child mortality (Bittles 1994; Majumdar et al. 1997; Dashtseren, 2002; Banerjee and Roy, 2002). Hence, the attainment of the best description of these interrelationships has remained a major and special goal of researchers from various disciplines.

The 1990's have seen a remarkable decrease in mortality among infants and children in most developing countries. Demographic and Health Survey contributed this to 5 factors, one of which is fertility behavior. Fertility is important in assessing the biological effects of consanguineous marriage on mortality. Fertility is greater in consanguineous marriages. This could be attributed to social factors such as younger age at marriage and first delivery. Studies revealed the impact of child mortality on fertility regulation behavior (Doepke, 2002), this can be explained by replacement and insurance effects. High fertility is a compensation for increased infant and childhood mortality as an attempt to achieve their desired family size (Rahman, 1998). Fertility control will be low among parents whose child dies, because they continue childbearing to replace, the child they lost. This is only if they didn't reach their desired family size.

Little is known about the possible linkage between parental consanguinity and offspring mortality in Egypt where it is widely practiced. Keeping this in mind, the present paper aims to examine genetic consequences of parental consanguinity on infant and child mortality in Egypt and its six major sub-divisions, namely: Upper Egypt Urban, Upper Egypt Rural, Lower Egypt Urban, Lower Egypt Rural, Urban Governorates, and Frontier Governorates, by controlling other important non-genetic proximate determinants of infant and child mortality.

Data and Methods

Our study is a secondary analysis of the data of 2000 Egypt Demographic Health Survey – a nationally representative sample of 16957 households from six administrative divisions of Egypt, namely, Frontier Governorate, Upper Egypt – Rural, Upper Egypt – Urban, Lower Egypt – Rural, Lower Egypt – Urban, and Urban Governorates. The sample from these six regions included 15573 ever-married women aged 15-49. The prevalence and pattern of consanguinity have been estimated from the answers to the two questions asked of all ever-married women in the Survey: “Prior to marriage, was your husband related to you in any way?” and if the answer to the first question was ‘yes’, “What type of relationship was it?” In order to assess the impact of consanguinity on offspring mortality the consanguineous group of women has been further divided into two separate groups, namely ‘close consanguinity’ and ‘remote consanguinity’ on the basis of the answers to the second question. Only first-degree relations have been included in the ‘close’; whereas second degree and other relations have been incorporated in the ‘remote’.

Two files were used: the file of women and the file of children <5. The survey provided detailed information on fertility, family planning, infant and child mortality, and maternal and child health care and nutrition. An index of living standards of the respondents has been calculated from information collected on the quality of housing, availability of electricity, sources of drinking water, nature of toilet facilities, fuel for cooking, ownership of goods etc. The index has been

divided into 3 categories based on the fact that the scores were normally distributed and thus one third of population is taken in each category as: low (0-15), medium (16-22), and high (23 and above). The Bivariate analysis revealed the effect of some background variables on mortality and consanguinity, such as education of the mother, birth interval, birth order, standard of living etc. Multivariate logistic regression models have been used considering two indices of mortality - infant mortality and child mortality, as dependent variable. The effect of the predictor on the dependent variable has been ascertained after controlling for other socioeconomic and non-genetic variables. Infant mortality has been calculated as proportion died within 0-11 months to births aged 12-59 months as denominator. Child mortality has been calculated as proportion died 12-59 months to births aged 5 years or above as denominator during the last 20 years prior to the survey. Finally, General Linear Model analysis is used to compare the mean number of children everborn between consanguineous marriages controlling for women's education, women's working status and Age of women at marriage. The Statistical Package for the Social Sciences (SPSS, 1988) is used for the analysis.

Results

Prevalence and Pattern of Consanguinity

Consanguineous marriages are still high in Egypt (38.9%). The prevalence of these marriages, however, varies by region. It ranges from 25.4% in Lower Egypt Urban to 55.2 % in Upper Egypt Rural (Table 1). The most common type of consanguineous marriages, in most regions, is that occurring between first cousins. Close consanguinity accounts for 22.2% of the total marriages; it is higher in rural areas and is found highest in Upper Egypt-Rural.

(Table 1 here)

Consanguinity and Offspring Mortality

Infant and Child Mortality are positively related to the degree of consanguinity. Looking at Infant and child mortality rates among consanguineous marriages have shown elevated mortality rates among the off-springs of close consanguineous marriages, where child and infant mortality are doubled among close consanguineous parents compared to parents having no relation (Figure 1).

(Figure 1 here)

Non-genetic Factors Affecting Offspring Mortality

The Bivariate analysis revealed that irrespective of the degree of consanguinity the non-genetic determinants, namely, standard of living, education of the

mother, region/residence, birth order, birth interval, age of the mother at delivery, size of the child, sex of the child, medical assistance during delivery and antenatal care has significant effect on offspring mortality (Table 2). An environmental factor that affects offsprings' mortality is standard way of living. An inverse relationship has been documented. In the population having low standard of living, child mortality among close consanguineous marriages is higher by 40% than non-consanguineous marriages and 6% than remote consanguineous marriages. In addition, off springs mortality varies by region/residence. It is very much true in case of Egypt where it is found significantly higher in Upper Egypt – Rural and lowest in Frontier Governorates.

Among other characteristics of the mother, maternal education has been regarded as the best predictor of offspring mortality. An inverse relationship has been found between the two: offsprings' mortality has been observed highest among illiterate mothers (Table 2). Irrespective of education, close consanguineous marriages had the highest mortality rates.

Another strong non-genetic determinants of infant and child mortality are the age of the mother at the time of birth, birth-interval, birth order and size of child at birth. Mothers delivering babies at the ages below 20 years carry greater risk of mortality in infancy and childhood of their newly born. For example, in the present sample, among close consanguineous marriages infant mortality is doubled for very young mothers as compared to mothers aged 20-29 years at the time of delivery. Among very young mothers, infant mortality rates were higher in close consanguineous marriages by 70%; were up by 27% in case of remote consanguineous marriages as compared to non-consanguineous unions. Similar observations were made in child mortality where rates were higher in close consanguineous marriages and remote consanguineous marriages by 59% and 28% than non-consanguineous, respectively. Further, children were at elevated risk of mortality if the interval between births was less than 24 months and it was higher for consanguineous marriages. Among the characteristics related to the child, children of birth order four and above had higher risk of offsprings' mortality. Similarly, this hazard was higher in case of consanguineous marriages. Also, child mortality was significantly higher in case of female children as compared to male children in close consanguineous marriages. Moreover, size of the child at birth had an inverse relationship with infant mortality. The smaller the child at birth the higher was the risk of mortality being approximately 5 times more among very small children as compared to having normal size deliveries. Again, among normal size children at birth, infant mortality rates were higher in close consanguineous marriages and remote consanguineous marriages by 63% and 6% as compared to non-consanguineous unions, respectively. The lack of antenatal care had an impact on infant mortality. Even if mothers had received antenatal care, infant mortality was 118% higher among close consanguineous marriages than that of parents with no relation (Table 2).

(Table 2 here)

Multivariate Analyses

Multivariate analyses have been carried out to find the effect of consanguinity on infant and child mortality after controlling the effects of non-genetic determinants. Two different logistic models have been used with two different indices of offspring mortality as dependent variables. Models I & II take infant mortality and child mortality as dependant variable, respectively. The results are presented in Table 3. As can be seen from the table, the risk of infant mortality was higher in case of consanguineous couples even after controlling for selected non-genetic predictors of infant mortality. The risk of mortality was 30% higher and was statistically significant in case of close consanguineous couples; it was higher by 19% among remote consanguineous couples but was not found significant. The results also show the impact of consanguinity on child mortality after controlling for other non-genetic factors. The risk of child mortality was higher among the close consanguineous couples by 50 % (odds ratio=1.52) and among remote consanguinity by 27% (odds ratio=1.27) as compared to no relation couples. The effects were found statistically significant.

(Table 3 here)

The General Linear Model Analysis revealed a higher mean number of children everborn among close consanguineous marriages (3.7), compared to no relation (3.2) and remote consanguineous marriages (3.6), after controlling for the effect of women's education, working status and age at marriage. This model is significant with a p-value less than 0.01 (Table 4).

(Table 4 here)

Discussion

Egypt has the largest population among 22 members of the League of Arab States. The prevalence of consanguinity varies between 20-50% in the region (Hamamy, 2003). It is widely practiced in Egypt (39%). However, the rate of consanguineous marriages in the country is lower than several other Arab countries. First cousin marriages constitute almost one third of all marriages in many Arab countries. The rates of these marriages differ between, as well as within, countries. In Egypt also, a wide variation in the prevalence of consanguineous marriages is observed in its six administrative divisions. Marriages between close relatives are found highest in Upper Egypt - Rural (30.8%); whereas they were lowest for Lower Egypt - Urban (14.5 %). The regional differentials in Infant and child mortality consistently reveal its pattern in close agreement with that of the prevalence of consanguinity in the six administrative divisions of Egypt. The results strongly support the close association of consanguinity with offsprings' mortality.

The influence of most of the non-genetic determinants of mortality is in the expected direction. Housing conditions and quality of sanitation are found significantly related with off-springs mortality. Economically weaker households were likely to experience higher infant and child mortality in the first five years of life as compared to the affluent households. Also, studies reveal that both infant mortality and child mortality are conditioned by the circumstances of childbirth and environmental factors to which mothers were exposed prior to giving birth. (Dashtseren, 2002). In addition, the inverse association between maternal education and offspring mortality found here is in agreement with the findings of many other studies, which show that maternal education decreases the risk of offspring mortality significantly (Hussein, 1998; Banerjee and Roy, 2002). This strong association between the two may be due to better personal hygiene, greater use of available health services and better child-care practices. Data from the Demographic and Health surveys indicate that in many Middle Eastern countries, infant and child mortality was higher for girls than that of boys. However, in case of the predictor 'utilization of health services', girls' mortality was lower as compared to boys (Hill, 1995). Contrary to general belief, the effect of 'medical assistance' during delivery is found inversely associated with the infant mortality. In the present study, higher infant mortality was observed where mothers received medical assistance as compared to those who did not at the time of delivery. This is in agreement with the findings of the study of Banerjee and Roy (2002) in India where it was found that births occurring in a hospital or clinic had higher odds of mortality compared with those delivered at home and without the help of any health professional. The authors argue that medical care is sought only in case of complications and unfortunately doctors were not always successful in saving lives of newborn (Banerjee and Roy, 2002).

Findings of the logistic regression analysis have shown that close consanguineous couples had 50% and 30% higher risks of death of their off-springs before their fifth birthday and first birthday, respectively than those couples who were not related by blood. In case of couples with remote consanguinity, the risk of dying of their offsprings before attaining fifth and first birthday was higher by 30% and 20%, respectively as compared to non-consanguineous couples.

As indicated in literature above fertility is higher in consanguineous marriages. A study by Frakenberg, on the relationship between Infant and child mortality and subsequent fertility in indonesia, showed that for certain groups of women, a child's death changes the fertility pattern of subsequent interval lengths relative to that pattern when a child survives (Frankenberg, 1991).

There are, however, two limitations of the present study: First, the information on congenital malformations, which are one of the leading causes of infant mortality particularly in the Arab region, was not available. This was a major impediment in assessing the effect in-depth of consanguinity on the mortality of

offsprings. Second, the probability of clustering of deaths in the same family may cause overestimation of infant and child mortality.

Conclusion

The findings of this study show the impact of consanguinity on infant and child mortality in Egypt, after controlling for their selected non-genetic determinants, leading indirectly to an increased fertility. Close consanguineous marriages, a cultural practice governed by consanguinity values and norms, increase a couple's risk of enduring the death of one or more of their children. Genetic disorders predictably will account for an increasing proportion of morbidity and mortality worldwide, and it is evident that this burden will fall disproportionately on countries and communities in which consanguinity is prevalent. Thus, an emphasis on the effect of consanguinity on mortality of off-springs, should be a part of intervention strategies that address the personal, emotional, and economic loss that families may face with the death of a child. A culturally appropriate approach for genetic counseling in relation to consanguineous marriage is required. Some Governments have put laws for premarital tests. The success of such counseling in Lebanon (Khlat et al. 1986) could be taken as an example. Public educational campaigns based on scientific evidence could help the purpose. In Egypt, where consanguinity is largely practiced, a proportion of offspring mortality may be prevented if society understands the importance of the issue and voluntarily decides to avoid marriage among biological relatives. However, it is important that the social and economic benefits of marriage to a close relative also be taken fully into consideration. Thus a national program should select strategies that have enough strength to dilute the cultural taboos linked with these social practices.

Holy Prophet Muhammad Mustafa (S.A.W.W) has said,

“.....don't get married to a very close relative because the offspring of such marriage will be weak. Defects out of such marriages will not be rectified until 3 generations of marriages of non-relatives”.

Reference

- Agwiser, A. (1990). Perinatal mortality at the armed forces hospital, Riyadh, Saudi Arabia: five year review of 22203 births. *Annals of Saudi medicine*, 10(3), 268-75.
- Al-Gazali, L., Bener, A., Abdulrazzaq, Y., Micallef, R., Khayyat, A., & Gaber, T. (1997). Consanguineous marriages in the United Arab Emirates. *Journal of biosocial Science*, 29 (4), 491-497.
- Banerjee, K., & Roy, TK. (2002). Parental consanguinity and offspring mortality: The search for possible linkage in the Indian context. *Asia-Pacific Population Journal*, 17 (1), 17-38.
- Bittles, A.H. (1994). The role and significance of consanguinity as a demographic variable. *Population and Development Review*, 20(3), 561-584.
- Bunday, S., & Alam, H. (1993). A five-year prospective study of the health of children in different ethnic groups, with particular reference to the effect of inbreeding. *European Journal of Human Genetics*, 1 (3), 206-219.
- Cabigon, J. (2002). Revisiting the 'Best' covariates of infant and child mortality: The Philippine case. Paper presented during the Bangkok Regional Population Conference "Southeast Asia's Population in a Changing Asian Context", 10-13 June, Bangkok, Thailand.
- Dashtseren, A. (2002). Determinants of infant and child mortality in Mongolia. Paper presented at the IUSSP Regional Conference, 9-13 June, Bangkok, Thailand.
- Doepke, M. (2002). Child mortality and fertility decline: Does the Barro-Becker Model fit the facts?. On-line working paper series. California Center for Population Research. University of California-Los Angeles.
- El-Hamzi, M., Al-Swailem, AR., Warsy, A., Al-Swailem, AM. & Sulaimani, R. (1995). Consanguinity among the Saudi Arabian population. *American Journal of medical Genetics*, 32, 623-626.
- Emery, A.E.H., Rimoin, D.L., eds. (1990). *Principles and practice of medical genetics*, Vols. 1 and 2, 2nd ed. Edinburgh, Churchill Livingstone.
- Federal Ministry Of Health, Sudan (1995). *Sudan Maternal and Child Health Survey*. Pan Arab Project for child development, Republic of Sudan and League of Arab States.
- Frankenberg, E. (1998). The relationship between infant and child mortality and subsequent fertility in Indonesia: 1971-1991. In: Montgomery, M., & Cohen, B.

(eds). *From death to birth: mortality decline and reproductive change*. Washington, D.C.: National Academy Press. 316-38.

Guo, G. (1993). Use of sibling data to estimate family mortality effects in Guatemala. *Demography* 30 (1), 15-32.

Hague, K., Bashi, O. (1988). Perinatal mortality at king khalid University Hospital, Riyadh. *Annals of Saudi medicine*, 8(3), 190-3

Hamamy, H. (2003). Consanguineous marriages in the Arab world. *The Ambassadors*. 6 (2),

HILL, K., & Upchurch, D. (1995). Gender differences in child health : Evidence from the Demographic and Health Surveys. *Population and development review*, 21(1), 127-151.

Hussain, R. (1998). The impact of consanguinity and inbreeding on perinatal mortality in Karachi, Pakistan. *Paediatrics Perinatal Epidemiology*, 12, 370-382.

Jurdi, R., & Saxena, P.C. (2003). The prevalence and correlates of consanguineous marriages in Yemen: Similarities and contrasts with other Arab countries. *Journal of biosocial Science*, 35 (1), 1-13.

Khoury, S., & Massad, D. (1992). Consanguineous marriages in Jordan. *American Journal of medical Genetics*, 43 (5), 769-775.

Khlat, M., Halabi, S., Khudr, A., & Der Kaloustian, V.M. (1986). Perception of consanguineous marriages and their genetic effects among a sample of couples from Beirut. *American Journal of medical Genetics*, 25, 299-306.

Ministry of Health, Bahrain (1991). *Health statistical abstract*.

Ministry Of Health, Kuwait (1996). *Kuwait Family Health Survey*. State of Kuwait and Gulf Family Health Survey.

Ministry of Health, United Arab Emirates (1992). *Statistical yearbook*.

Ministry of Public Health, Qatar (1993). *Vital statistics annual report 1992*.

Ministry Of Public Health, Tunisia (1996). *Tunisian Maternal and Child Health Survey*. Pan Project for Child Development, Republic of Tunisia and League of Arab States.

National Population Council, Egypt (1996). *Egypt Demographic and Health Survey*. Arab Republic of Egypt and Macro International Inc., Calverton, Maryland.

Rahman, M. (1998). The effect of child mortality on fertility regulation in rural Bangladesh. *Studies in Family Planning*, 29(3), 268-281.

Shah, G., Toney, M., & Pitcher, B. (1998). Consanguinity and child mortality: The risk faced by families. *Population Research and Policy Review*, 17, 275-283.

Stern, C. (1949). *Principles of Human Genetics* (San Francisco, California, W.H. Freedman and Company).

SPSS. 1988. Statistical Package for the Social Sciences. *SPSS-X: introductory statistics guide for release 3*. Chicago, IL: SPSS, Inc.

Temtamy, S.A., Abdel Meguid, N., Mazen, I., Ismail, S.R., Kassem N.S., & Bassiouni R. (1998). A genetic epidemiological study of malformations at birth in Egypt. *EMHJ*, 4 (2), 252-259

World Health Organization Regional Office for the Eastern Mediterranean (1993). *Data provided by the Health Situation and Trend Assessment Unit*.

Table 1: Distribution of Ever-married Women, Aged 15-49 Years, by their Marriage Pattern, Egypt and its Governorates, 2000.


Region of Egypt *	Type of Relationship With the Husband						Total No. of Ever-married Women	
	No relation		Close Consanguinity*		Remote Consanguinity**		N	%
	N	%	N	%	N	%		
Urban Governorates	2212	71.3	519	16.7	371	12.0	3102	100
Lower Egypt - Urban	1367	74.7	265	14.5	199	10.9	1831	100
Lower Egypt - Rural	2706	633.3	989	23.1	582	13.6	4277	100
Upper Egypt - Urban	1040	62.3	316	18.9	314	18.8	1670	100
Upper Egypt - Rural	1678	44.8	1151	30.8	914	24.4	3743	100
Frontier Governorates	509	53.6	212	22.3	228	24.0	949	100
Total	9512	61.1	3452	22.2	2608	16.7	15572	100

Note: Egypt is divided into 26 governorates. Four Urban (Cairo, Alexandria, Port Said, and Suez) with no rural population, while the other 22 have both urban and rural. 9 of the mixed governorates are in the Nile Delta (Lower Egypt), and 8 in the Nile Valley (Upper Egypt), while the remaining 5 Frontier Governorates are on the eastern and western boundaries of Egypt.

* Includes marriage among first degree relations.

** Comprises marriage among second degree and other relations.

Figure 1: Infant and child mortality rates by consanguinity, Egypt, 2000.



Note: Infant mortality was calculated as proportion died within 0-11 months to births aged 12-59 months as denominator during 4 years prior to the survey, using the file of children < 5. Child mortality was calculated as proportion died 12-59 months to births aged 5 years or above as denominator during the last 20 years prior to the survey, using birth history from the women's file. Both rates were per 1000 livebirths.

Table 3: Logistic regression coefficients giving extent of effects of selected determinants of Infant and child mortality, Egypt, 2000.

Variable	Infant mortality (N=8978)		Child mortality (N= 30754)	
	OR	95% CI	OR	95% CI
Consanguinity				
No relation	1		1	
Close	1.295†	1.028-1.632	1.516*	1.294-1.776
Remote	1.188	0.911-1.550	1.269†	1.051-1.533
Standard of living				
Low	1		1	
Medium	1.112	0.883-1.400	0.925	0.791-1.081
High	0.811	0.589-1.118	0.486*	0.368-0.642
Region				
Urban Governorates	1		1	
Lower Egypt Urban	0.965	0.618-1.506	1.361	0.966-1.918
Lower Egypt Rural	0.904	0.634-1.289	1.054	0.794-1.398
Upper Egypt Urban	1.239	0.831-1.847	1.352	0.971-1.883
Upper Egypt Rural	1.226	0.861-1.747	1.506*	1.139-1.992
Frontier Governorates	0.748	0.296-1.893	0.783	0.358-1.714
Maternal education				
Illiterate	1		1	
Primary	1.023	0.777-1.347	0.834†	0.696-1.000
Secondary & Higher	0.993	0.761-1.297	0.376*	0.273-0.516
Age of mother at birth				
<20	1.701*	1.213-2.351	1.279†	1.020-1.604
20-29	1		1	
30-49	1.243	0.964-1.603	1.116	0.936-1.331
Birth Interval				
<24	1		1	
24-47	0.617*	0.485-0.785	0.435*	0.374-0.506
48+	0.351*	0.245-0.502	0.289*	0.225-0.371
Birth Order				
1	0.670*	0.498-0.901	1.034	0.823-1.299
2-3	1		1	
4+	1.328†	1.014-1.740	1.409*	1.181-1.680
Sex of the child				
Male	1		1	
Female	0.880	0.724-1.069	1.233*	1.073-1.416
Size of the child				
Normal	1		---	---
Small	2.434*	1.900-3.119	---	---
Very small	5.031*	3.640-6.954	---	---
Assistance during delivery				
Some assistance	1		---	---
No one	0.752	0.298-1.903	---	---
Antenatal care				
Yes	1		---	---
No	1.276†	1.027-1.584	---	---

Note: --- refers to variables that were not included in the model.

* p-value < 0.01. † p-value < 0.05

Table 2: Infant and child mortality rates by selected determinants, Egypt, 2000.

Variable	Infant mortality			Child mortality		
	No	Close	Remote	No	Close	Remote
Standard of living						
Low	56	63	73	35	49	46
Medium	53	78	58	25	42	28
High	28	65	49	5	20	19
Region						
Urban Governorates	34	73	39	12	24	9
Lower Egypt Urban	37	63	32	17	39	16
Lower Egypt Rural	38	63	52	20	37	32
Upper Egypt Urban	56	88	34	18	23	46
Upper Egypt Rural	71	68	88	39	54	41
Frontier Governorates	27	59	33	10	22	24
Maternal education						
Illiterate	52	70	84	32	47	42
Primary	64	63	72	19	39	34
Secondary & Higher	37	68	32	6	14	6
Age of mother at birth						
<20	69	117	92	29	46	36
20-29	41	58	43	20	36	32
30-49	51	72	90	24	48	31
Birth Interval						
<24	73	90	111	39	71	55
24-47	44	65	52	15	31	25
48+	26	38	47	12	12	14
Birth Order						
1	41	78	43	17	26	17
2-3	41	51	57	16	38	32
4+	60	80	80	33	49	41
Sex of the child						
Male	50	67	65	21	38	26
Female	42	69	58	23	43	40
Size of the child						
Normal	32	52	49	---	---	---
Small	106	99	113	---	---	---
Very small	194	285	263	---	---	---
Assistance during delivery						
Some assistance	45	66	61	---	---	---
No one	29	97	59	---	---	---
Antenatal care						
Yes	33	72	46	---	---	---
No	60	62	75	---	---	---

Note: Infant and child mortality rates were calculated per 1000 livebirths.

--- Information on Antenatal care, Assistance during delivery and size of the child was collected for children born during the four years prior to the survey.

Table 4: General Linear Model Analysis of the mean number of children everborn by consanguinity controlling for women's education, working status and Age at marriage, Egypt, 2000.

Source of variation	Mean	SD	N	P-value
Consanguinity				
No relation	3.2	2.1	8555	
Close	3.7	2.2	2985	< 0.01
Remote	3.6	2.3	2225	
Covariates *				
Women's education				< 0.01
Working status				< 0.01
Age at marriage				< 0.01

Note: the women's file is used in the analysis taking women as the unit of analysis.

* Women's education is categorical (Illiterate, Primary, Secondary and Higher), Working status (working and not working), Age at marriage is continuous.